

Invariant submanifolds of LCS-manifold admitting semi-symmetric metric connection

¹Ananda K, ²Sudhir Patel,

^{1,2}Assistant Professor, New Horizon College of Engineering, Bangalore, India,

¹anandak.nhce@gmail.com, ²sudhirfriends@gmail.com

Abstract: In this paper, we study invariant submanifolds of (LCS)-manifold admitting semi-symmetric metric connection. We prove that the submanifolds also carry the semi-symmetric metric connection. Further, we also consider recurrent, 2-recurrent and generalized 2-recurrent submanifolds of (LCS)-manifold admitting semi-symmetric metric connection and we investigate the conditions for all the above mentioned submanifolds to be totally geodesic.

Key Words: Invariant submanifold, (LCS)-manifold, recurrent, 2-recurrent, generalized 2-recurrent and totally geodesic.

I. INTRODUCTION

Let (M, g) be an n -dimensional Riemannian manifold with metric g and Let ∇ be the Levi-civita connection on M . A linear connection $\bar{\nabla}$ on (M, g) is said to be Semi-symmetric [1] if the torsion tensor T of connection $\bar{\nabla}$ satisfies

$$T(X, Y) = W(X)Y - W(Y)X$$

where W is a 1-form on M . Further let ρ be the vector field associated with it, that is $W(X) = g(X, \rho)$ for any differentiable vector field X on M . A Semi-symmetric connection $\bar{\nabla}$ is called Semi-symmetric metric connection [2] if it satisfies $\bar{\nabla}g = 0$. In 1924, Friedman and Schouten [1] introduced the idea of semi-symmetric linear connection on a differentiable manifold. Further in 1932, The idea of metric connection with torsion on a Riemannian manifold was introduced by H.A. Hayden [2]. Later, Yano [3] synthesized the notion of semi-symmetric connection and a metric connection with torsion on a Riemannian manifold in 1970. After the properties of semi-symmetric metric connection have studied by many authors like K.S. Amur and S.S. Pujar [4], C.S. Bagewadi, D.G. Prakasha and Venkatesha [5,6], A. Sharfuddin and S.I. Hussain[7], U.C. De and G. Pathak [8] etc. Recently, in 2003 A.A. Shaikh [9] introduced and studied Lorentzian concircular structure manifold (briefly (LCS)-manifold) which generalizes the notion of LP-sasakian manifolds, introduced by Matsumoto [10] in 1989. Then Shiakh and Baishya [11,12] have been studied on the applications of $(LCS)_n$ - manifolds to the general theory of relativity and cosmology. It is to be noted that $(LCS)_n$ manifold remains invariant under a D-homothetic transformation [13]. On the other hand in 2008, A.A. Shaikh, T.Basu and S. Eyasmin [20]

proved that existence of ϕ -recurrent $(LCS)_3$ manifold which is neither locally symmetric nor locally ϕ -symmetric by non trivial examples. Later, G.T. Shreenivasa, Venkatesha, C.S. Bagewadi [23] also studied $(LCS)_{2n+1}$ -manifolds in 2009. Further, in 2011 Atceken and Hui have been studied on the submanifold of an $(LCS)_n$. The (LCS) manifolds have been also studied by Atceken [14], Narain and Yadav [15], Prakasha [16], Shaikh [17], Shaikh et al. [14,18], Shaikh and Bihn [21], shaikh and Hui [21], Sreenivasa et al. [22], Yadav et al. [23] and others. Recently, in 2016 Shaikh, Matsuyama and Hui have been studied on invariant submanifolds of $(LCS)_n$ -manifolds and it deals with the study of some basic properties of invariant submanifolds of $(LCS)_n$ -manifolds.

If $\bar{\nabla}$ denotes semi-symmetric metric connection on a contact metric manifold, then it is given by [13]

$$\bar{\nabla}_X Y = \nabla_X Y + \eta(Y)X - g(X, Y)\xi, \quad (1.1)$$

Where $\eta(Y) = g(Y, \xi)$.

The covariant differential of the p^{th} order, $p \geq 1$, of a $(0, k)$ -tensor field T , $k \geq 1$, defined on a Riemannian manifold (M, g) with the Levi-civita connection ∇ , is denoted by $\nabla^p T$.

The tensor T is said to be recurrent if it satisfies

$$(\nabla T)(X_1, \dots, X_K; X)T(Y_1, \dots, Y_K) = (\nabla T)(Y_1, \dots, Y_K; X)T(X_1, \dots, X_K), \quad (1.2)$$

On M . where $X, Y, X_1, Y_1, \dots, X_K, Y_K \in TM$. From (1.2) it follows that at a point $x \in M$, if the tensor T is non zero, then there exist a unique 1-form ϕ and a $(0, 2)$ -tensor ψ , defined on a neighbourhood U of x such that

$$\nabla T = T \otimes \phi, \quad \phi = d(\log \|T\|). \quad (1.3)$$

Similarly, the tensor T is said to be 2-recurrent [13] if it satisfies

$$(\nabla^2 T)(X_1, \dots, X_K; X, Y)T(Y_1, \dots, Y_K) = (\nabla^2 T)(Y_1, \dots, Y_K; X, Y)T(X_1, \dots, X_K), \quad (1.4)$$

On M . Where $X, Y, X_1, Y_1, \dots, X_K, Y_K \in TM$. Now from (1.4) it follows that at a point $x \in M$, if the tensor T is non zero, then there exist a $(0, 2)$ -tensor ψ , defined on a neighborhood U of x such that

$$\nabla^2 T = T \otimes \psi, \quad (1.5)$$

hold on U , where $\|T\|$ denotes the norm of T and $\|T\|^2 = g(T, T)$.

The tensor T is said to be generalized 2-recurrent if

$$\begin{aligned} & ((\nabla^2 T)(X_1, \dots, X_K; X, Y) \\ & - (\nabla T \otimes \phi)(X_1, \dots, X_K; X, Y))T(Y_1, \dots, Y_K) = \\ & ((\nabla^2 T)(Y_1, \dots, Y_K; X, Y) - \\ & (\nabla T \otimes \phi)(Y_1, \dots, Y_K; X, Y))T(X_1, \dots, X_K), \end{aligned} \quad (1.6)$$

hold on M , where ϕ is a 1-form on M . From (1.6) it follows that at a point $x \in M$ if the tensor T is non zero, then there exists a $(0, 2)$ -tensor ψ , defined on a neighborhood U of x such that

$$\nabla^2 T = \nabla T \otimes \phi + T \otimes \psi, \quad (1.7)$$

holds on U .

II. ISOMETRIC IMMERSION

Let $f : (M, g) \rightarrow (\tilde{M}, \tilde{g})$ be an isometric immersion from an n -dimensional Riemannian manifold (M, g) into $(n+d)$ -dimensional Riemannian manifold (\tilde{M}, \tilde{g}) , $n \geq 2$, $d \geq 1$. We denote

by ∇ and $\tilde{\nabla}$ as Levi-Civita connection of M^n and \tilde{M}^{n+d} respectively. Then the formulas of Gauss and Weingarten are given by

$$\tilde{\nabla}_X Y = \nabla_X Y + \sigma(X, Y), \quad (2.1)$$

$$\tilde{\nabla}_X N = -A_N X + \nabla_X^\perp N, \quad (2.2)$$

for any tangent vector fields X, Y and the normal vector field N on M . Where

σ -second fundamental form,

A -the shape operator,

∇^\perp -is the normal connection.

If the second fundamental form σ is identically zero, then the manifold is said to be totally geodesic. The second fundamental form σ and A_N are related by

$$\tilde{g}(\sigma(X, Y), N) = g(A_N X, Y), \quad (2.3)$$

for tangent vector fields X, Y .

The first and second covariant derivatives of the second fundamental form σ are given by

$$(\tilde{\nabla}_X \sigma)(Y, Z) = \nabla_X^\perp(\sigma(Y, Z)) - \sigma(\nabla_X Y, Z) - \sigma(Y, \nabla_X Z), \quad (2.4)$$

$$(\tilde{\nabla}^2 \sigma)(Z, W, X, Y) = (\tilde{\nabla}_X \tilde{\nabla}_Y \sigma)(Z, W),$$

$$= \nabla_X^\perp((\tilde{\nabla}_Y \sigma)(Z, W)) - (\tilde{\nabla}_Y \sigma)(\nabla_X Z, W) -$$

$$(\tilde{\nabla}_X \sigma)(Z, \nabla_Y W) - (\tilde{\nabla}_{\nabla_X Y} \sigma)(Z, W), \quad (2.5)$$

respectively, where $\tilde{\nabla}$ is called the Vander Waerden-Bortolotti connection of M [7].

If

$$\tilde{\nabla} \sigma = 0$$

then M is said to have parallel second fundamental form [7].

III. LCS-MANIFOLD

In 2003 A. A Shaikh the first author introduced the notion of Lorentzian concircular structure manifold (briefly, $(LCS)_n$ -manifolds) and these are applied in General theory of relativity and cosmology.

An n -dimensional Lorentzian Manifold M is a smooth connected para-compact Hausdorff manifold with a Lorentzian metric g . that is, M admits a smooth symmetric tensor field g of type $(0, 2)$ such that for each point $p \in M$, the tensor $g_p: T_p M \times T_p M \rightarrow \mathbb{R}$ is a non-degenerate inner product of signature $(-, +, \dots, +)$, where $T_p M$ denotes the tangent vector space of M at p and \mathbb{R} is the real number space. A non-zero vector $v \in T_p M$ is said to be timelike if it satisfies $g_p(v, v) < 0$. The category to which a given vector fields is called its casual character.

In a Lorentzian manifold (M, g) , a vector field P defined by

$$g(X, P) = A(X),$$

for any X on M , is said to be a concircular vector field if

$$(\nabla_X A)(Y) = \alpha\{g(X, Y) + \omega(X)A(Y)\}, \quad (3.1)$$

for Y on M , where α is a non-zero scalar and ω is a closed 1-form.

Let M be Lorentzian manifold admitting a unit timelike concircular vector field ξ , called the structure vector field of the manifold. Then we have

$$g(\xi, \xi) = -1, \quad (3.2)$$

Since ξ is a unit concircular vector field, it follows that there exist a non-zero 1-form η such that

$$g(X, \xi) = \eta(X), \quad (3.3)$$

The following equation $(\nabla_X \eta)(Y) = \alpha\{g(X, Y) + \eta(X)\eta(Y)\}$,

$$(3.4)$$

holds for all vector fields X, Y on M and α is a non-zero scalar function satisfies

$$\nabla_X \alpha = X(\alpha) = d\alpha(X) = \rho\eta(X), \quad (3.5)$$

ρ being a certain scalar function given by

$$\rho = -\xi(\alpha), \quad (3.6)$$

If we put

$$\nabla_X \xi = \alpha \varphi X, \quad (3.7)$$

Then from (3.4) and (3.7) we have

$$\varphi X = X + \eta(X)\xi, \quad (3.8)$$

From which it follows that

$$\varphi^2 = X + \eta(X)\xi, \quad (3.9)$$

that is, φ is a symmetric (1,1) tensor field, called the structure tensor of the manifold. The n -dimensional Lorentzian manifold M together with the unit timelike concircular vector field ξ , its associated 1-form η , and an (1, 1) tensor field φ is said to be a Lorentzian concircular structure manifold (briefly, $(LCS)_n$ -manifold). Especially, if $\alpha=1$, then we can obtain the LP-Sasakian structure of Matsumoto.

In $(LCS)_n$ -manifold, the following results hold

$$\eta(\xi) = g(\xi, \xi) = -1, \phi^2 X = X + \eta(X)\xi, g(X, \xi) = \eta(X), \phi\xi = 0, \eta\phi = 0. \quad (3.10)$$

$$g(\phi X, \phi Y) = g(X, Y) + \eta(X)\eta(Y), \quad (3.11)$$

$$\eta(R(X, Y)Z) = (\alpha^2 - \rho)\{g(Y, Z)\eta(X) - g(X, Z)\eta(Y)\}, \quad (3.12)$$

$$S(X, \xi) = (\alpha^2 - \rho)(n - 1)\eta(X), \quad (3.13)$$

for any vector fields X, Y, Z on M and $\alpha^2 - \rho \neq 0$, where R -curvature tensor and S -Ricci tensor of the manifold.

In an $(LCS)_n$ -manifold, we also have the following results

$$(\nabla_X \eta)(Y) = (\nabla_Y \eta)(X) \quad (3.14)$$

$$d\eta(X, Y) = 0 \quad (3.15)$$

We also mention that, in an $(LCS)_n$ -manifold the symmetric (1, 1) tensor field ϕ is idempotent and hence the eigen value of ϕ is either 1 or 0.

IV. INVARIANT SUBMANIFOLDS OF (LCS)-MANIFOLD ADMITTING SEMI-SYMMETRIC METRIC CONNECTION

If \tilde{M} is a (LCS) -manifold with structure tensors $(\bar{\phi}, \bar{\xi}, \bar{\eta}, \bar{g})$ then we know that its invariant submanifold M has the induced (LCS) -structure (ϕ, ξ, η, g) .

A submanifold M of a (LCS) -manifold \tilde{M} with a semi-symmetric metric connection is called an invariant submanifold of \tilde{M} with a semi-symmetric connection, if for each $x \in M, \phi(T_x M) \subset T_x M$. The study of such submanifolds for the different type of contact

manifolds have been carried out by authors of [4, 12], [26]-, [29], [33, 37]

As a consequence, ξ becomes tangent to M . For an invariant submanifold of a (LCS) manifold with a semi-symmetric metric connection, We have

$$\sigma(X, \xi) = 0, \quad (4.1)$$

for any vector X tangent to M .

Let \bar{M} be a (LCS) -manifold admitting a semi-symmetric metric connection $\bar{\nabla}$.

Lemma 4.1. *Let M be an invariant submanifold of contact metric manifold \tilde{M} which admits semi-symmetric connection $\bar{\nabla}$ and let σ and $\bar{\sigma}$ be the second fundamental form with respect to Levi-Civita connection and semi-symmetric metric connection then*

- (a) M admits semi-symmetric metric connection,
- (b) the second fundamental form with respect to $\bar{\nabla}$ and $\bar{\bar{\nabla}}$ are equal.

Proof. : we know that the contact metric structure $(\bar{\phi}, \bar{\xi}, \bar{\eta}, \bar{g})$ on \tilde{M} induces (ϕ, ξ, η, g) on invariant submanifold. By virtue of (1.1), we get

$$\bar{\nabla}_X Y = \bar{\nabla}_X Y + \eta(Y)X - g(X, Y)\xi. \quad (4.2)$$

By using (2.1) in (4.2)

$$\bar{\nabla}_X Y = \nabla_X Y + \sigma(X, Y) + \eta(Y)X - g(X, Y)\xi. \quad (4.3)$$

Now gauss formula (2.1) with respect to semi-symmetric metric connection is given by

$$\bar{\nabla}_X Y = \bar{\nabla}_X Y + \bar{\sigma}(X, Y). \quad (4.4)$$

Equating (4.3) and (4.4), we get (1.1)

$$\bar{\sigma}(X, Y) = \sigma(X, Y). \quad (4.5)$$

V. RECURRENT INVARIANT SUBMANIFOLDS OF (LCS)-MANIFOLD ADMITTING SEMI-SYMMETRIC METRIC CONNECTION

We consider invariant submanifolds of a (LCS) -manifold When σ is recurrent, 2-recurrent, generalized 2-recurrent and M has parallel third fundamental form with respect to semi-symmetric metric connection. We write the equations (2.4) and (2.5) with respect to semi-symmetric metric connection in the form

$$\begin{aligned} (\bar{\nabla}_X \sigma)(Y, Z) &= \bar{\nabla}_X^\perp(\sigma(Y, Z)) - \sigma(\bar{\nabla}_X Y, Z) - \sigma(Y, \bar{\nabla}_X Z), \quad (5.1) \\ (\bar{\nabla}_X^2 \sigma)(Z, W, X, Y) &= (\bar{\nabla}_X \bar{\nabla}_Y \sigma)(Z, W) = \bar{\nabla}_X^\perp((\bar{\nabla}_Y \sigma)(Z, W)) - (\bar{\nabla}_Y \sigma)(\bar{\nabla}_X Z, W) - (\bar{\nabla}_X \sigma)(Z, \bar{\nabla}_Y W) - (\bar{\nabla}_{\bar{\nabla}_X Y} \sigma)(Z, W). \quad (5.2) \end{aligned}$$

Theorem 5.1. *Let M be an invariant submanifold of $(LCS)_n$ -manifold \tilde{M} admitting semi-symmetric metric connection. Then σ is recurrent with respect to semi-symmetric metric connection if and only if it is totally geodesic with respect to Levi-Civita connection.*

Proof. :Let σ be recurrent with respect to semi-symmetric metric connection. Then from (1.3) we get

$$(\bar{\nabla}_X \sigma)(Y, Z) = \phi(X) \sigma(Y, Z),$$

Where ϕ is a 1-form on M . By using (5.1) and put $Z = \xi$ in the above equation, we have

$$\bar{\nabla}_X^\perp(\sigma(Y, \xi) - \sigma(\bar{\nabla}_X Y, \xi) - \sigma(Y, \bar{\nabla}_X \xi)) = \phi(X) \sigma(Y, \xi),$$

Which by virtue of (4.1) reduces to

$$-\sigma(\bar{\nabla}_X Y, \xi) - \sigma(Y, \bar{\nabla}_X \xi) = 0. \quad (5.4)$$

Using (1.1), (3.7), (3.10) and (4.1) in (5.4)

$$-\alpha \sigma(Y, \phi X) + \sigma(Y, X) = 0. \quad (5.5)$$

Now replace X by ϕX and by virtue of (3.10), (4.1) in (5.5), we get

$$(1 - \alpha) \sigma(Y, X) = 0. \quad (5.6)$$

We get $\sigma(X, Y) = 0$, if $\alpha \neq 1$ Therefore M is totally geodesic provided $\alpha \neq 1$

Theorem 5.2. Let M be an invariant submanifold of a (LCS)-manifold \tilde{M} admitting semi-symmetric metric connection. Then M has parallel third fundamental form with respect to semi-symmetric metric connection if and only if it is totally geodesic with respect to Levi-Civita connection.

Proof. Let M has parallel third fundamental form with respect to semi-symmetric metric connection. then we have

$$(\bar{\nabla}_X \bar{\nabla}_Y \sigma)(Z, W) = 0$$

Taking $W = \xi$ and using (5.2) in the above equation, we have

$$\bar{\nabla}_X^\perp \left((\bar{\nabla}_Y \sigma)(Z, \xi) - (\bar{\nabla}_Y \sigma)(\bar{\nabla}_X Z, \xi) - (\bar{\nabla}_X \sigma)(Z, \bar{\nabla}_Y \xi) - (\bar{\nabla}_{\bar{\nabla}_X Y} \sigma)(Z, \xi) \right) = 0. \quad (5.7)$$

$$0 = \bar{\nabla}_X^\perp \{ \sigma(\bar{\nabla}_Y Z, \xi) + \sigma(Z, \bar{\nabla}_Y \xi) \} - \bar{\nabla}_Y^\perp \{ \sigma(\bar{\nabla}_X Z, \xi) + \sigma(\bar{\nabla}_Y \bar{\nabla}_X Z, \xi) + 2\sigma(\bar{\nabla}_X Z, \bar{\nabla}_Y \xi) - \bar{\nabla}_X^\perp \{ \sigma(Z, \bar{\nabla}_Y \xi) + \sigma(Z, \bar{\nabla}_X \bar{\nabla}_Y \xi) + \sigma(\bar{\nabla}_{\bar{\nabla}_X Y} \xi) \}. \quad (5.8)$$

In view of (1.1), (3.7), (3.10) and (4.1) the above result (5.8) gives

$$0 = -2\bar{\nabla}_X^\perp \sigma(Z, \alpha \phi Y) + 2\bar{\nabla}_X^\perp \sigma(Z, Y) + \sigma(Z, \nabla_X \alpha \phi Y) - \sigma(Z, \nabla_X \eta(Y) \xi) + \alpha \sigma(Z, \phi \nabla_X Y) + \alpha \eta(Y) \sigma(Z, \phi X) - \alpha \sigma(Z, \nabla_X Y) - \eta(Y) \sigma(Z, X) + 2\alpha \sigma(\nabla_X Z, \phi Y) - 2\sigma(\nabla_X Z, Y) + 2\alpha \eta(Z) \sigma(X, \phi Y) - 2\eta(Z) \sigma(X, Y). \quad (5.9)$$

Put $Y = \xi$ and using (3.7), (3.10) and (4.1) in (5.9), we get

$$0 = (\alpha^2 + 1) \sigma(Z, X) - 2\alpha \sigma(Z, \phi X). \quad (5.10)$$

Multiply by $(\alpha^2 + 1)$ in (5.10), we get

$$0 = (\alpha^2 + 1)^2 \sigma(Z, X) - 2\alpha(\alpha^2 + 1) \sigma(Z, \phi X). \quad (5.11)$$

Put $X = \phi X$ in (5.10), we get

$$0 = (\alpha^2 + 1) \sigma(Z, \phi X) - 2\alpha \sigma(Z, X). \quad (5.12)$$

Multiply by 2α in (5.12), we get

$$0 = 2\alpha(\alpha^2 + 1) \sigma(Z, \phi X) - 4\alpha^2 \sigma(Z, X). \quad (5.13)$$

Adding equation (5.11) and (5.13), we get $\alpha \neq \pm 1$ and $\sigma(X, Z) = 0$. Thus M is totally geodesic. The converse statement is trivial. This proves the theorem

Corollary 5.1. :Let M be an invariant submanifold of a (LCS)-manifold \tilde{M} admitting semi-symmetric metric connection. Then σ is 2-recurrent with respect to semi-symmetric metric connection if and only if it is totally geodesic with respect to Levi-Civita connection.

Proof. Let σ be 2-recurrent with respect to semi-symmetric metric connection. From (1.5), we have

$$(\bar{\nabla}_X \bar{\nabla}_Y \sigma)(Z, W) = \sigma(Z, W) \phi(X, Y).$$

Taking $W = \xi$ and using (5.2) in the above equation, we have

$$\bar{\nabla}_X^\perp \left((\bar{\nabla}_Y \sigma)(Z, \xi) - (\bar{\nabla}_Y \sigma)(\bar{\nabla}_X Z, \xi) - (\bar{\nabla}_X \sigma)(Z, \bar{\nabla}_Y \xi) - (\bar{\nabla}_{\bar{\nabla}_X Y} \sigma)(Z, \xi) \right) = \sigma(Z, \xi) \phi(X, Y). \quad (5.14)$$

In view of (4.1) and (5.1) we write (5.14) in the form

$$0 = \bar{\nabla}_X^\perp \{ \sigma(\bar{\nabla}_Y Z, \xi) + \sigma(Z, \bar{\nabla}_Y \xi) \} - \bar{\nabla}_Y^\perp \{ \sigma(\bar{\nabla}_X Z, \xi) + \sigma(\bar{\nabla}_Y \bar{\nabla}_X Z, \xi) + 2\sigma(\bar{\nabla}_X Z, \bar{\nabla}_Y \xi) - \bar{\nabla}_X^\perp \{ \sigma(Z, \bar{\nabla}_Y \xi) + \sigma(Z, \bar{\nabla}_X \bar{\nabla}_Y \xi) + \sigma(\bar{\nabla}_{\bar{\nabla}_X Y} Z, \xi) + \sigma(Z, \bar{\nabla}_{\bar{\nabla}_X Y} \xi) \}. \quad (5.15)$$

Using (1.1), (3.7), (3.10) and (4.1) in (5.15), we get

$$0 = -2\bar{\nabla}_X^\perp \sigma(Z, \alpha \phi Y) + 2\bar{\nabla}_X^\perp \sigma(Z, Y) + \sigma(Z, \nabla_X \alpha \phi Y) - \sigma(Z, \nabla_X \eta(Y) \xi) + \alpha \sigma(Z, \phi \nabla_X Y) + \alpha \eta(Y) \sigma(Z, \phi X) - \alpha \sigma(Z, \nabla_X Y) - \eta(Y) \sigma(Z, X) + 2\alpha \sigma(\nabla_X Z, \phi Y) - 2\sigma(\nabla_X Z, Y) + 2\alpha \eta(Z) \sigma(X, \phi Y) - 2\eta(Z) \sigma(X, Y). \quad (5.16)$$

Put $Y = \xi$ and using (3.7), (3.10) and (4.1) in (5.16), we get

$$0 = (\alpha^2 + 1) \sigma(Z, X) - 2\alpha \sigma(Z, \phi X) \quad (5.17)$$

Multiply by $(\alpha^2 + 1)$ in (5.17), we get

$$0 = (\alpha^2 + 1)^2 \sigma(Z, X) - 2\alpha(\alpha^2 + 1) \sigma(Z, \phi X) \quad (5.18)$$

Put $X = \phi X$ in (5.17), we get

$$0 = (\alpha^2 + 1) \sigma(Z, \phi X) - 2\alpha \sigma(Z, X) \quad (5.19)$$

Multiply by 2α in (5.19), we get

$$0 = 2\alpha(\alpha^2 + 1) \sigma(Z, \phi X) - 4\alpha^2 \sigma(Z, X) \quad (5.20)$$

Adding equation (5.18) and (5.20), we get $\alpha \neq \pm 1$ and $\sigma(X, Z) = 0$. Thus M is totally geodesic. The converse statement is trivial. This proves the theorem.

Theorem 5.3. :Let M be an invariant submanifold of a (LCS)-manifold \tilde{M} admitting semi-symmetric metric connection. Then σ is generalised 2-recurrent with respect to semi-symmetric metric connection if and only if it is totally geodesic with respect to Levi-Civita connection.

Proof. Let σ be a generalized 2-recurrent with respect to semi-symmetric metric connection.

From (1.7), We have

$$(\bar{\nabla}_X \bar{\nabla}_Y \sigma)(Z, W) = \psi(X, Y) \sigma(Z, W) + \phi(X) (\bar{\nabla}_Y \sigma)(Z, W). \quad (5.21)$$

Where ψ and ϕ are 2-recurrent and 1-form respectively.

Taking $W = \xi$ in (5.21) and using (4.1), We get

$$(\bar{\nabla}_X \bar{\nabla}_Y \sigma)(Z, \xi) = \phi(X) (\bar{\nabla}_Y \sigma)(Z, \xi).$$

Using (4.1), (5.1) and (5.2) in above equation, We get

$$\begin{aligned} \bar{\nabla}_X^{\perp} \left((\bar{\nabla}_Y \sigma)(Z, \xi) \right) - (\bar{\nabla}_Y \sigma)(\bar{\nabla}_X Z, \xi) - (\bar{\nabla}_X \sigma)(Z, \bar{\nabla}_Y \xi) \\ - (\bar{\nabla}_{\bar{\nabla}_X Y} \sigma)(Z, \xi) = \\ - \phi(X) \{ \sigma(\bar{\nabla}_Y Z, \xi) + \sigma(Z, \bar{\nabla}_Y \xi) \} \end{aligned} \quad (5.22)$$

By using (4.1) and by the virtue of (5.1), We write (5.22) in the form

$$\begin{aligned} \bar{\nabla}_X^{\perp} \{ \sigma(\bar{\nabla}_Y Z, \xi) + \sigma(Z, \bar{\nabla}_Y \xi) \} - \bar{\nabla}_Y^{\perp} \{ \sigma(\bar{\nabla}_X Z, \xi) \} + \\ \sigma(\bar{\nabla}_Y \bar{\nabla}_X Z, \xi) + 2\sigma(\bar{\nabla}_X Z, \bar{\nabla}_Y \xi) - \bar{\nabla}_X^{\perp} \{ \sigma(Z, \bar{\nabla}_Y \xi) \} + \\ \sigma(Z, \bar{\nabla}_X \bar{\nabla}_Y \xi) + \sigma(\bar{\nabla}_{\bar{\nabla}_X Y} Z, \xi) + \sigma(Z, \bar{\nabla}_{\bar{\nabla}_X Y} \xi) = \\ - \phi(X) \{ \sigma(\bar{\nabla}_Y Z, \xi) + \sigma(Z, \bar{\nabla}_Y \xi) \}. \end{aligned} \quad (5.23)$$

In view of (1.1), (3.7), (3.10) and (4.1) in (5.23), We get

$$\begin{aligned} -2\bar{\nabla}_X^{\perp} \sigma(Z, \alpha\phi Y) + 2\bar{\nabla}_X^{\perp} \sigma(Z, Y) + \sigma(Z, \nabla_X \alpha\phi Y) - \\ \sigma(Z, \nabla_X \eta(Y)\xi) + \alpha\sigma(Z, \phi \nabla_X Y) + \alpha\eta(Y)\sigma(Z, \phi X) - \\ \alpha\sigma(Z, \nabla_X Y) - \eta(Y)\sigma(Z, X) + 2\alpha\sigma(\nabla_X Z, \phi Y) - \\ 2\sigma(\nabla_X Z, Y) + 2\alpha\eta(Z)\sigma(X, \phi Y) - 2\eta(Z)\sigma(X, Y) = \\ - \phi(X) \{ \sigma(Z, \alpha\phi Y) - \sigma(Z, Y) \}. \end{aligned} \quad (5.24)$$

Put $Y = \xi$ and using (3.7), (3.10) and (4.1) in (5.24), We get

$$0 = (\alpha^2 + 1)\sigma(Z, X) - 2\alpha\sigma(Z, \phi X). \quad (5.25)$$

Multiply by $(\alpha^2 + 1)$ in (5.25), we get

$$0 = (\alpha^2 + 1)^2 \sigma(Z, X) - 2\alpha(\alpha^2 + 1)\sigma(Z, \phi X). \quad (5.26)$$

Put $X = \phi X$ in (5.25), we get

$$0 = (\alpha^2 + 1)\sigma(Z, \phi X) - 2\alpha\sigma(Z, X). \quad (5.27)$$

Multiply by 2α in (5.27), we get

$$0 = 2\alpha(\alpha^2 + 1)\sigma(Z, \phi X) - 4\alpha^2\sigma(Z, X). \quad (5.28)$$

Adding equation (5.26) and (5.28), we get $\alpha \neq \pm 1$ and $\sigma(X, Z) = 0$. Thus M is totally geodesic. The converse statement is trivial. This proves the theorem.

VI. CONCLUSION

From the Theorems (5.1), (5.2) and (5.3) and from the Corollary (5.1), we conclude that:

If M is an invariant submanifolds of LCS manifold admitting semi-symmetric metric connection then the following conditions are equivalent:

- i. The submanifold M is totally geodesic with respect to the Levi-Civita connection;
- ii. Second fundamental form σ is recurrent with respect to semi-symmetric metric connection;
- iii. The submanifold M has parallel third fundamental form with respect to semi-symmetric metric connection;
- iv. Second fundamental form σ is 2-recurrent with respect to semi-symmetric metric connection;
- v. Second fundamental form σ is generalized 2-recurrent with respect to semi-symmetric metric connection.

REFERENCES

- [1] A. Friedmann and J.A. Schouten, Uber die geometric der halbsymmetrischen Ubertragung, math. Zeitschr:21 (1924), 211-223.
- [2] H.A.Hayden, Subspaces of space with torsion, Proc. London Math. Soc. 34 (1932), 27-50.
- [3] K. Yano, On semi-symmetric metric connection, Resv. Roumaine, Math. Press Apple.15 (1970), 1579- 1586.
- [4] K.S. Amur and S.S. Pujar, On submanifolds of a Riemannian manifold admitting a metric semi-symmetric connection, Tensor; N.S. 32 (1978), 35-38.
- [5] C.S. Bagewadi, On totally real submanifolds of a Kahlerian manifold admitting semi-symmetric metric F-connection, Indian J. Pure Appl. Math. 13 (5) (1982), 528-536.
- [6] C.S. Bagewadi, D.G. Prakasha and Venkatesha, Projective curvature tensor on a kenmotsu manifold with respect to semi-symmetric metric connection, Stud. Cercet Stiint. Ser: Mat. Univ. Bacau. 17 (2007), 21-32.
- [7] A. Sharafuddin and S.I. Hussain, Semi-symmetric metric connection in almost contact manifolds, Tensors; N.S.30 (1976), 133-139.
- [8] U.C. De and G. Pathak, On a semi-symmetric metric connection in Kenmotsu manifolds, Bull. Calcutta Math. Soc. 94 (40) (2002), 319-324.

- [9] A.A. Shaikh, On Lorentzian almost paracontact manifolds with a structure of the concircular type, *kyung-pook. Math. J.*43 (2003) 305-314.
- [10] K. Matsumoto, On Lorentzian almost paracontact manifolds, *Bull. Yamagata Univ. Nat. Sci.*12 (1989) 151-156.
- [11] I. Mihai, R. Rosca, On Lorentzian para-Sasakian manifolds, World Scientific Publication, Singapore (1992), pp. 155-169.
- [12] D. Nirmla, M.S. Siddesha and C.S. Bagewadi, On Invariant submanifolds of Lorentzian β -Kenmotsu manifold, *Asian Journal of Mathematics and Computer Research*, 19(4) (2017), 203-213.
- [13] A.A. Shaikh, K.K. Baishya, On Concircular structure spacetimes, *J. Math. Stat.* 1 (2005),129-132.
- [14] A.A. Shaikh, K.K. Baishya, On Concircular structure spacetimes, *N, Am. J. Appl. Sci.*3 (4) (2006), 1790-1794.
- [15] A.A. Shaikh, H. Ahmed, Some Transformations of (LCS) n -manifolds, *Tsukuba J. Math.* 38 (2014), 1-24.
- [16] M. Atceken, On Geometry of submanifolds of (LCS) n -manifolds, *Int. J. Math. Math. Sci.* (2012),doi:10.1155/2012/304647.
- [17] D. Narain, S. Yadav, On weak concircular symmetries of (LCS) $2n+1$ -manifolds, *Glob. J. Sci. Front. Res.* 12 (2012), 85-94.
- [18] D.G. Prakasha, On Riccy η -recurrent (LCS) n -manifolds, *Acta Univ. Apulensis.* 24 (2010), 109-118.
- [19] A.A. Shaikh, Some results on (LCS) n -manifolds *J. Korean Math. Soc.* 46 (2009), 449-461.
- [20] A.A. Shaikh, T. Basu, S.Eyasmin, On locally φ -symmetric (LCS) n -manifolds, *Int. J. Pure Appl. Math.* (8) (2007), 1161-1170.
- [21] A.A. Shaikh, T. Basu, S.Eyasmin, On existence of φ -recurrent (LCS) n -manifolds, *Extracta Math.* 23 (2008), 71-83.
- [22] A.A. Shaikh, T.Q. Binh, On weakly symmetric (LCS) n -manifolds, *J. Adv. Math. Stud.* 2 (2009), 75-90.
- [23] A.A. Shaikh, S.K. Hui, On generalized φ -recurrent (LCS) n -manifolds, in: proceedings of the AIP conference, 1309, 2010, pp 419-429
- [24] G.T. Sreenivasa, Venkatesha, C.S. Bagewadi, Some results on (LCS) $2n+1$ -manifolds, *Bull. Math. Anal. Appl.* 1 (3) (2009), 64-70.
- [25] S.K. Yadav, P.K. Dwivedi, D. Suthar, On (lcs) $2n+1$ -manifolds satisfying certain consitions on the concircular curvature tensor, *Thai J. Math.* 9 (3) (2011), 597-603.
- [26] M.S. Siddesha and C.S. Bagewadi, Submanifold of a (k, μ) -Contact manifold, *CUBO A mathematical Journal*, 18(01) (2016), 59-68.
- [27] M.S. Siddesha and C.S. Bagewadi, On some classes of invariant submanifolds (k, μ) -contact manifold, *Journal of Informatics and Mathematical Sciences*, 9(1) (2017), 13-26.
- [28] M.S. Siddesha and C.S. Bagewadi, Invariant submanifolds (k, μ) -contact manifold admitting quarter sym- metric metric connection, *International Journal of Mathematics Trends and Technology (IJMTT)*, 34 (2)- June 2016, 48-53.
- [29] S. Sular and C. Ozgur, On some submanifolds of Kenmotsu manifolds, *Chaos, Solitons and Fractals.* 42 (2009), 1990-1995.
- [30] Z.I. Szabo, Structure theorems on Riemannian manifolds satisfying $R(X, Y) \cdot R = 0$, I Local version. *J. Differ. Geom.* 17 (1982), 531-582.
- [31] Z.I. Szabo, Structure theorems on Riemannian manifolds satisfying $R(X, Y) \cdot R = 0$, II Global version. *Geom. Dedicata* 19 (1985), 65-108.
- [32] S. Tanno, Ricci curvatures of contact Riemannian manifolds, *Tohoku Mathemat- ical Journal* 40, 441 (1988).
- [33] M.M. Tripathi, T. Sasahara and J.S. Kim, On invariant submanifolds of contact metric manifolds, *Tsukuba J. Math.* 29(2) (2005), 495-510.
- [34] L. Verstraelen, Comments on pseudosymmetry in the sense of Ryszard Deszcz, In *Geometry and Topology of submanifolds of World Scientific Publishing*, River Edge vol. VI (1994), 199-209.
- [35] K. Yano, Concircular geometry I. concircular transformations in:, *Proceedings of the Imperial Academy Tokyo*, 16 (1940), 195-200.
- [36] K. Yano and M. Kon, Structures on manifolds, World scientific publishing, (1984).
- [37] Ahmet Yildz and Cengizhan Murathan, Invariant submanifolds of Sasakian space forms, *J. Geom.* 95 (2009), 135-150.